



## Early Journal Content on JSTOR, Free to Anyone in the World

This article is one of nearly 500,000 scholarly works digitized and made freely available to everyone in the world by JSTOR.

Known as the Early Journal Content, this set of works include research articles, news, letters, and other writings published in more than 200 of the oldest leading academic journals. The works date from the mid-seventeenth to the early twentieth centuries.

We encourage people to read and share the Early Journal Content openly and to tell others that this resource exists. People may post this content online or redistribute in any way for non-commercial purposes.

Read more about Early Journal Content at <http://about.jstor.org/participate-jstor/individuals/early-journal-content>.

JSTOR is a digital library of academic journals, books, and primary source objects. JSTOR helps people discover, use, and build upon a wide range of content through a powerful research and teaching platform, and preserves this content for future generations. JSTOR is part of ITHAKA, a not-for-profit organization that also includes Ithaka S+R and Portico. For more information about JSTOR, please contact [support@jstor.org](mailto:support@jstor.org).

*Reports of States and yearly and monthly reports of cities.*

ALABAMA—*Mobile*.—Month of November, 1896. Estimated population, 37,817. Total deaths, 73, including phthisis pulmonalis, 13; enteric fever, 3; diphtheria, 1, and croup, 2.

CONNECTICUT—*Westport*.—Month of January, 1897. Estimated population, 4,000. Total deaths, 4. No deaths from contagious diseases.

FLORIDA.—Month of September, 1896. Reports to the State board of health from 45 counties, having an aggregate population of 391,418, show a total of 298 deaths, including smallpox, 6; enteric fever, 14; phthisis pulmonalis, 16; croup, 1; measles, 2, and whooping cough, 1.

Month of October, 1896. Total deaths, 304, including smallpox, 1; phthisis pulmonalis, 27; enteric fever, 18; measles, 1, and whooping cough, 3.

*Pensacola*.—Month of January, 1897. Estimated population, 15,000. Total deaths, 29, including 6 from phthisis pulmonalis.

GEORGIA—*Columbus*.—Month of January, 1897. Estimated population—white, 10,276; colored, 9,027; total, 19,303. Deaths—white, 7; colored, 11; total 18, including phthisis pulmonalis, 1, and croup, 1.

KENTUCKY—*Louisville*.—Month of January, 1897. Estimated population, 211,100. Total deaths, 261, including phthisis pulmonalis, 29; enteric fever, 7; scarlet fever, 1; diphtheria, 2, and croup, 3.

MICHIGAN.—Week ended January 30, 1897. Reports to the State board of health from 62 observers indicate that erysipelas decreased in area of prevalence. Phthisis pulmonalis was reported present during the week at 176 places, measles at 49, diphtheria at 47, scarlet fever at 29, enteric fever at 24, and whooping cough at 18 places.

OHIO—*Dayton*.—Month of January, 1897. Estimated population, 85,000. Total deaths, 88, including phthisis pulmonalis, 15; enteric fever, 1; diphtheria, 6; scarlet fever, 1, and whooping cough, 1.

VIRGINIA—*Richmond*.—Month of January, 1897. Estimated population, 103,560. Total deaths, 130, including phthisis pulmonalis, 9; diphtheria, 1, and whooping cough, 1.